



Return to Learn After Concussion

ECHO IDAHO: K12 School Nurses

January 8th, 2025

*Kristi Pardue, MS, CCC-SLP, CBIS
Manager, St. Luke's Sports Medicine Concussion Clinic*



Please keep in mind that your School District policies and Health Services procedures, medication administration protocols, process guidelines, remain the guiding principles to your practice. None of the planners or presenters for this educational activity have relevant financial relationship(s) to disclose with ineligible companies whose primary business is producing, marketing, selling, re-selling, or distributing healthcare products used by or on patients.

Learning Objectives

- Describe recent updates in the world of concussion. Attention to the 6th International Conference on Concussion in Sport.
- Understand the importance of a school concussion management plan.
- Identify three resources for school nurses when working with students during concussion recovery.




Concussion Definition

- A concussion is a traumatic brain injury that can't be seen on routine x-rays, CT scans or MRIs. It affects the way a person may think and remember things and can cause a variety of symptoms. Any blow to the head, face or neck, or a blow to the body that jars your head, could cause a concussion.
- *No abnormality is seen on standard structural neuroimaging studies.*
- May involve LOC, but most patients do not have a LOC.
- Symptoms and signs may be present immediately or evolve over minutes or hours or a day and commonly resolve within days but may be prolonged.

Consensus statement

Consensus statement on concussion in sport: the 6th International Conference on Concussion in Sport—Amsterdam, October 2022





COGNITIVE

Feeling Mentally Foggy
Difficulty Concentrating
Difficulty Remembering
Repeats Questions
Feeling Mentally Slowed Down
Forgetful of Recent Information
Confused About Recent Events
Answers Questions Slowly

PHYSICAL

Headache
Nausea/Vomiting
Balance Problems
Numbness/Tingling
Sensitivity to Light/Noise
Visual Problems
Dizziness
Dazed or Stunned

Irritability
Sadness
More Emotional
Nervousness

EMOTIONAL

Drowsiness
Sleeping Less Than Usual
Sleeping More Than Usual
Trouble Falling Asleep

SLEEP



Legislation

Idaho's Youth Athlete Concussion Legislation HB 557/Section 33-1625

Primary Tenets of Idaho's Legislation

- ✓ Education coaches, parents & athletes
- ✓ Obtain written consent from parent/guardian
- ✓ Remove athlete from play
- ✓ Return to learn before return to play
- ✓ Obtain medical clearance from a qualified healthcare providers knowledgeable about concussion before return to play

Applies to Middle School, Junior High School, and High School sanctioned sports

****Youth sports organizations are strongly encouraged to comply with this legislation and afforded liability protection if comply**



<https://legislature.idaho.gov/statutesrules/idstat/Title33/T33CH16/SECT33-1625/>

Legislation

Idaho Youth Athlete Concussion Legislation HB 557/Section 33-1625

- *Return to learn before return to play*


“Students who have sustained a concussion and return to school may need informal or formal accommodations, modifications of curriculum, and monitoring by medical or academic staff until the student is fully recovered. A student athlete should be able to resume all normally scheduled academic activities without restrictions or the need for accommodation prior to receiving authorization to return to play by a qualified healthcare professional as defined in subsection (6) of this section.”



What's the Latest?

Consensus statement

Consensus statement on concussion in sport: the 6th International Conference on Concussion in Sport—Amsterdam, October 2022

Jon S Patricios ¹, Kathryn J Schneider ², Jiri Dvorak ³,
Osman Hassan Ahmed ^{4,5}, Cheri Blauwet ^{6,7}, Robert C Cantu,^{8,9}
Gavin A Davis ^{10,11}, Ruben J Echemendia ^{12,13}, Michael Makdissi,^{14,15}
Michael McNamee,^{16,17} Steven Broglio ¹⁸, Carolyn A Emery ²,
Nina Feddermann-Demont,^{19,20} Gordon Ward Fuller ²¹, Christopher C Giza,^{22,23}
Kevin M Guskiewicz,²⁴ Brian Hainline ²⁵, Grant L Iverson ^{26,27},
Jeffrey S Kutcher,²⁸ John J Leddy ²⁹, David Maddocks,³⁰ Geoff Manley ³¹,
Michael McCrea ³², Laura K Purcell,³³ Margot Putukian ³⁴, Haruhiko Sato ³⁵,
Markku P Tuominen,³⁶ Michael Turner ^{37,38}, Keith Owen Yeates ³⁹,
Stanley A Herring,^{40,41} Willem Meeuwisse⁴²

For numbered affiliations see end of article.

Correspondence to

Dr Kathryn J Schneider, Sport Injury Prevention Research Centre, Faculty of Kinesiology, University of Calgary, Calgary, AB T2N 1N4, Canada; kjschnei@ucalgary.ca

ABSTRACT

For over two decades, the Concussion in Sport Group has held meetings and developed five international statements on concussion in sport. This 6th statement summarises the processes and outcomes of the 6th International Conference on Concussion in Sport held in Amsterdam on 27–30 October 2022 and should be read in conjunction with the (1) methodology paper that outlines the consensus process in detail and (2)

methodology. The purpose of this Statement is to provide a summary of the evidence and practice recommendations based on science and expert panel consensus recommendations at the time of the conference. Additional outputs of the consensus process include freely available evidence-informed tools to assist in the detection and assessment of SRC, including the Concussion Recognition Tool-6 (CRT6), Sport Concussion Assessment Tool-6

Br J Sports Med: first published as 10.1136/bjsports-2023-106998 on 14 June 2023. D



Key Points

Strict rest is not beneficial, may be harmful

Early return to physical activity recommended

Reduced (not eliminated) screen time

Aerobic exercise may be advanced based on tolerance, or per exercise testing



Updated Tools

CRT6™

Concussion Recognition Tool

To Help Identify Concussion in Children, Adolescents and Adults

What is the Concussion Recognition Tool?

A concussion is a brain injury. The Concussion Recognition Tool 6 (CRT6) is to be used by non-medically trained individuals for the identification and immediate management of suspected concussion. It is not designed to diagnose concussion.

Recognise and Remove

Red Flags: CALL AN AMBULANCE

If ANY of the following signs are observed or complaints are reported after an impact to the head or body the athlete should be immediately removed from play/game/activity and transported for urgent medical care by a healthcare professional (HCP):

<ul style="list-style-type: none"> • Neck pain or tenderness • Seizure, 'fits', or convulsion • Loss of vision or double vision • Loss of consciousness • Increased confusion or deteriorating conscious state (becoming less responsive, drowsy) 	<ul style="list-style-type: none"> • Weakness or numbness/tingling in more than one arm or leg • Repeated Vomiting • Severe or increasing headache • Increasingly restless, agitated or combative • Visible deformity of the skull
--	---

Remember

- In all cases, the basic principles of first aid should be followed: assess danger at the scene, check airway, breathing, circulation, look for reduced awareness of surroundings or slowness or difficulty answering questions.
- Do not attempt to move the athlete (other than required for airway support) unless trained to do so.
- Do not remove helmet (if present) or other equipment.
- Assume a possible spinal cord injury in all cases of head injury.
- Athletes with known physical or developmental disabilities should have a lower threshold for removal from play.

This tool may be freely copied in its current form for distribution to individuals, teams, groups, and organizations. Any alteration (including translations and digital re-formatting), re-branding, or sale for commercial gain is not permissible without the expressed written consent of BMJ.

Developed by: The Concussion in Sport Group (CISG)

Supported by:

Br J Sports Med: first published as 10.1136/bjsports-2023-107021 on 14 June 2023. Downloaded from <http://open.bmj.com/> on January 6, 2025 by guest. Protected by copyright.

CRT6

Concussion Recognition Tool

To Help Identify Concussion in Children, Adolescents and Adults

1: Visible Clues of Suspected Concussion

Visible clues that suggest concussion include:

- Loss of consciousness or responsiveness
- Lying motionless on the playing surface
- Falling unprotected to the playing surface
- Disorientation or confusion, staring or limited responsiveness, or an inability to respond appropriately to questions
- Dazed, blank, or vacant look
- Seizure, fits, or convulsions
- Slow to get up after a direct or indirect hit to the head
- Unsteady on feet / balance problems or falling over / poor coordination / wobbly
- Facial injury

2: Symptoms of Suspected Concussion

Physical Symptoms	Changes in Emotions
Headache	More emotional
"Pressure in head"	More irritable
Balance problems	Sadness
Nausea or vomiting	Nervous or anxious
Drowsiness	
Dizziness	
Blurred vision	
More sensitive to light	
More sensitive to noise	
Fatigue or low energy	
"Don't feel right"	
Neck Pain	

Changes in Thinking
Difficulty concentrating
Difficulty remembering
Feeling slowed down
Feeling like "in a fog"

Remember, symptoms may develop over minutes or hours following a head injury.

3: Awareness

(Modify each question appropriately for each sport and age of athlete)

Failure to answer any of these questions correctly may suggest a concussion:

- "Where are we today?"
- "What event were you doing?"
- "Who scored last in this game?"
- "What team did you play last week/game?"
- "Did your team win the last game?"

Any athlete with a suspected concussion should be - IMMEDIATELY REMOVED FROM PRACTICE OR PLAY and should NOT RETURN TO ANY ACTIVITY WITH RISK OF HEAD CONTACT, FALL OR COLLISION, including SPORT ACTIVITY until ASSESSED MEDICALLY, even if the symptoms resolve.

Athletes with suspected concussion should **NOT**:

- Be left alone initially (at least for the first 3 hours). Worsening of symptoms should lead to immediate medical attention.
- Be sent home by themselves. They need to be with a responsible adult.
- Drink alcohol, use recreational drugs or drugs not prescribed by their HCP
- Drive a motor vehicle until cleared to do so by a healthcare professional

British Journal of Sports Medicine

Echemendia RI, et al. Br J Sports Med June 2023 Vol 57 No 11

Br J Sports Med: first published as 10.1136/bjsports-2023-107021 on 14 June 2023. Downloaded from <http://open.bmj.com/> on January 6, 2025 by guest. Protected by copyright.



Living Guideline Return to School/Learn Protocol

Step	Activity	Examples of activities
1	Activities of daily living and relative rest* (Maximum of 24-48 hours)	Activities at home such as social interactions and light walking that do not result in more than mild and brief** exacerbation (worsening) of concussion symptoms. Minimize screentime.
2	School activities with encouragement to return to school as soon as possible (as tolerated)	Reading or other cognitive activities at school or at home. Goal: Increase tolerance to cognitive work, and connect socially with peers. Take breaks and adapt activities if concussion symptom exacerbation (worsening) is more than mild and brief**. Clearance from your doctor is not required to return to low-risk in-person or at-home school activities. A complete absence from the school environment for more than one week is not generally recommended.
3	Part-time or full days at school with academic accommodations if needed	Gradual reintroduction of school work. May require partial school days with access to breaks throughout the day, or with academic accommodations to tolerate the classroom or school environment. Gradually reduce accommodations and increase workload until full days without concussion-related accommodations are tolerated.
4	Return to school full-time. No academic accommodations (related to concussion)	Return to full days at school and academic activities without requiring concussion-related accommodations. Medical clearance is NOT required to return to school.

Date Updated: Sept 2023

Instructions:

Students should begin a gradual increase in their cognitive load with the goal of minimizing time away from the school environment. The return to school should not be restricted if the student is tolerating full days. Progression through the strategy may be slowed when there is more than a mild and brief symptom exacerbation**; however, missing more than one week of school is not generally recommended.

Definitions:

- *Relative rest: activities of daily living including walking and other light physical and cognitive activities are permitted as tolerated.
- **Mild exacerbation (worsening) of symptoms: No more than a 2-point increase when compared with the pre-activity value on a 0-10-point symptom severity scale***. *Brief* exacerbation of symptoms: Worsening of symptoms for up to 1 hour.
- ***0-10 point symptom severity scale: Please see the [Visual analog scale](#) for an example of a 0-10 symptom severity scale.

These definitions and instructions were harmonized with and modified with permission from the [Amsterdam International Consensus Statement on Concussion in Sport](#)

Living Guideline Return to Activity/Sport Protocol

Step	Activity	Examples of activities
1	Activities of daily living and relative rest* (Maximum of 24-48 hours)	Activities at home such as social interactions and light walking that do not result in more than mild and brief** exacerbation (worsening) of concussion symptoms. Minimize screentime.
2	Aerobic exercise Step 2A: Light effort (up to approx 55% of maximum heart rate) Step 2B: Moderate effort (up to approx 70% of maximum heart rate)	Start with stationary cycling or walking at slow to medium pace. Take a break and modify activities as needed with the aim of gradually increasing tolerance and the intensity of aerobic activities. Light resistance training that does not result in more than mild and brief** exacerbation (worsening) of concussion symptoms. Goal: increase the heart rate.
3	Individual sport-specific activities that do not have a risk of inadvertent head impact	Sport-specific training away from the team sport environment (e.g., running, change of direction, and/or individual training drills and individual gym class activities that do not have a risk of head impact and are supervised by a teacher or coach). Goal: Increase the intensity of aerobic activities and introduce low-risk sport-specific movements and changing of directions.
Medical clearance and a full return to school are required to progress to Step 4		
4	Non-contact training drills and activities	Exercise to high intensity including more challenging training drills and activities (e.g., passing drills, multiplayer training, high-intensity exercises, supervised non-contact gym class activities, and practices without body contact). Goal: Resume usual intensity of exercise, coordination, and activity-related cognitive skills
5	Return to all non-competitive activities, all gym class activities, and full-contact practices	Participate in higher-risk activities including normal training activities, all school gym-class activities, and full-contact sports practices and scrimmages. Avoid competitive gameplay. Goal: return to activities that have a risk of falling or body contact, restore game-play confidence, and have coaches assess functional skills.
6	Return to sport	Normal, unrestricted competitive gameplay, school gym class, and physical activities

This return-to-activity/sport table was modified with permission from the [Amsterdam International Consensus Statement on Concussion in Sport](#)

Instructions:

Begin Step 1 (i.e., relative rest) within 24 hours of injury, with progression through each subsequent step taking a minimum of 24 hours. If more than mild exacerbation (worsening) of symptoms (i.e., more than 2 points on a 0-10 scale***) occurs during Steps 1-3, stop the activity and attempt to exercise the next day. People experiencing concussion-related symptoms during Steps 4-6 should return to Step 3 to establish full resolution of symptoms with exertion before engaging in at-risk activities. Written determination of medical clearance should be provided before unrestricted Return to Sport as directed by local laws and/or sporting regulations.

Definitions:

- *Relative rest: activities of daily living including walking and other light physical and cognitive activities are permitted as tolerated.
- **Mild exacerbation (worsening) of symptoms: No more than a 2-point increase when compared with the pre-activity value on a 0-10-point symptom severity scale***. *Brief* exacerbation of symptoms: Worsening of symptoms for up to 1 hour.
- ***0-10 point symptom severity scale: Please see the [Visual analog scale](#) for an example of a 0-10 symptom severity scale. These definitions were harmonized with and modified with permission from the [Amsterdam International Consensus Statement on Concussion in Sport](#)



Return to School

This tool is a guideline for managing a student's return to school following a concussion and does not replace medical advice. Every concussion is unique, and recovery is very different for each individual. Not everyone will require a Return to School Strategy. Timelines and activities may vary based on direction from a doctor, nurse practitioner, or licensed healthcare professional with relevant training. Note: For information about returning to activities that pose risk of head contact, please refer to Return to School or Return to Activity.

STEP 1:	STEP 2:	STEP 3:	STEP 4:
<p>Activities of daily living and relative rest*</p> <ul style="list-style-type: none"> Maximum of 24-48 hours Activities at home such as social interactions and light walking that do not result in more than mild and brief** exacerbation (worsening) of concussion symptoms. <p>Examples:</p> <ul style="list-style-type: none"> Preparing meals Housework Light walking <ul style="list-style-type: none"> Minimize screen time for the first 24-48 hours following concussion. Avoid driving during the first 24-48 hours after a concussion. <p>Contact school to create a Return to School plan.</p>	<p>School activities (as tolerated)</p> <ul style="list-style-type: none"> Returning to school as soon as possible (as tolerated) is encouraged. Reading or other cognitive activities at school or at home. Goal: Increase tolerance to cognitive work, and connect socially with peers. Take breaks and adapt activities if concussion symptom exacerbation (worsening) is more than mild and brief.** Use of devices with screens may be gradually resumed, as tolerated. Clearance from your doctor is not required to return to low-risk in-person or at-home school activities. A complete absence from the school environment for more than one week is not generally recommended. <p>Communicate with school on student's progression</p>	<p>Part-time or full-time days at school with accommodations (if needed)</p> <ul style="list-style-type: none"> Gradually reintroduce schoolwork. May require accommodations, such as: <ul style="list-style-type: none"> Partial school days with access to breaks throughout the day Academic accommodations (extra time to complete work, reduced workload) to tolerate the classroom or school environment. <p>Communicate with school on student's progression.</p>	<p>Return to school full-time</p> <p>Return to full days at school and academic activities without requiring accommodation (related to the concussion).</p> <p>Note: Medical clearance is NOT required to return to school</p> <p>For returning to P.E. or sports, please refer to Return to Sport protocol.</p>
<p>Activities of daily living, as tolerated</p> <p>After a maximum of 24-48 hours after injury, BEGIN STEP 2</p>	<p>Return to school as soon as possible, as tolerated</p> <p>If can tolerate school activities, BEGIN STEP 3</p>	<p>Gradually reduce accommodations and increase workload</p> <p>If can't tolerate full days without concussion-related accommodations, BEGIN STEP 4</p>	<p>Full academic load (no academic accommodations related to the concussion)</p> <p>Return to School completed</p>

Students should begin a gradual increase in their cognitive load with the goal of minimizing time away from the school environment. The return to school should not be restricted if the student is tolerating full days. Progression through the strategy may be slowed when there is more than a mild and brief symptom exacerbation***; however, missing more than one week of school is not generally recommended. Driving should resume after consultation with a doctor, nurse practitioner, or healthcare professional.

*Relative rest: activities of daily living including walking and other light physical and cognitive activities are permitted as tolerated.

Mild exacerbation (worsening) of symptoms: No more than a 2-point increase when compared with the pre-activity value on a 0-10-point symptom severity scale.*Brief* exacerbation of symptoms: Worsening of symptoms for up to 1 hour.

***0-10 point symptom severity scale: Please see the [Visual Analog Scale](#) for an example of a 0-10 symptom severity scale.

www.cattonline.com

Adapted from: Zemek, R., Reed, N., Dawson, J., et al. "Living Guideline for Pediatric Concussion Care." www.pedsconcussion.com
© BCIRPU. All rights reserved | Version 14: Updated December 2024

©ATT° | CONCUSSION AWARENESS TRAINING TOOL

BC INJURY research and prevention unit
www.injuryresearch.bc.ca



Case Example

DOI 10/1/24 due to MVA – 11th grade student

Thought they were okay and it would go away but symptoms worsened, and school nurse recommended seeing a doctor.

10/24/24 went to urgent care who recommended they go to the ED. Seen at ED on 10/24/24 (didn't go directly to ED – went home instead) and dx with concussion and instructed to follow-up with PCP.

Concussion Clinic received referral on 11/12/24 from PCP.

As of 11/4/24 was “tolerating school a half day once a week” “getting sent home every time they try to attend if they had a headache” now concerned about being behind in school.

Saw Concussion Clinic physician on 11/18/24. Physician asked for assistance to work with school & family and provided Academic Recovery Recommendations to family to give to school.



Case Example Cont....

- Call with parent and school revealed:

- ✓ *Communication had been with attendance line not school nurse*
- ✓ *School nurse reported they had received no medical documents or indication as to why student had been missing school. Had not heard from student after initially recommending they follow-up with a physician.*
- ✓ *School team very willing to help student*
- ✓ *Initial medical team had given no guidance regarding return to learn (urgent care, ED, PCP)*

Common themes regardless of patient:

- ✓ *Communication between school, parent, student, and healthcare team breakdowns*
- ✓ *Need for concussion management plan (based on current research)*
- ✓ *One or two teachers would benefit from understanding concussion*
- ✓ *School team sending student home when they became symptomatic*



Benefits of a Concussion Management Plan

- Key team members identified.
- School team knows what to do and why and the why behind it before it happens.
- Not just about symptoms – “what can I do to help you?” Likely different for each student.
- Much easier to follow a plan in place vs create each time.
- Focus on positive recovery from concussion vs impairment.
- School staff knowledge on how to support student with adjustments and why vs “when are you going to be over this???”
- Will be ready to move to more formal supports if symptoms persist, however, most of the time if a plan is in place, this will be avoided.



Initial Considerations

****Positive expectations with appropriate accommodations/adjustments*

- Look at student's schedule to determine easiest transition
 - Avoid cafeteria, identify alternative location with 1-2 friends
 - Exempt from non-essential work
 - No testing or no more than one a day
 - Focus on core classes initially depending on time of year
 - Consider grade and time of year
 - Limit homework
 - State and standardized tests delayed until fully tolerating school
 - Class transition before bell
 - Rest breaks – identify location and schedule of break time
 - Grades are not an indicator in high achieving students
 - Consider premorbid difficulties
 - Prior history of difficulties does not negate the possibility of new difficulties
- *“Screen time” can mean something different to each person. i.e. 6 hours of video games vs texting a friend. Might be tiring but its not dangerous.*




Resources

- stlukesonline.org/concussion

Updated IHSAA Statewide Concussion Education Video
Fall 2024

Concussion Education and Training Video Share



When in Doubt, Sit Them Out

The CDC estimates that 1.8 to 3.6 million sports- and activity-related concussions happen each year in the United States. Because so many people are affected, providers at St. Luke's have established St. Luke's Sports Medicine Concussion Clinic to support patients and their primary care physicians in southern Idaho. Our mission is to set the standard in the comprehensive care of concussion through a multidisciplinary approach to prevention and care, as well as research, family support, and community reintegration.



Concussion



St. Luke's Sports Medicine Concussion Clinic

The CDC estimates that 1.8 to 3.6 million sports- and activity-related concussions happen each year in the United States. Because so many people are affected, providers at St. Luke's have established St. Luke's Sports Medicine Concussion Clinic to support patients and their primary care physicians in southern Idaho. Our mission is to set the standard in the comprehensive care of concussion through a multidisciplinary approach to prevention and care, as well as research, family support, and community reintegration.


To make an appointment at St. Luke's Sports Medicine Concussion Clinic, please contact us:

- Phone: (208) 381-2665
- Fax: (208) 381-9201 for referring providers

Concussion Clinic Services Concussion Clinic Specialties Concussion Outreach and Education

8th grade health curriculum video – Boise School District

Concussion Discussion for Students Share




The Boise School District & St. Luke's Sports Medicine Concussion Clinic are partnering with the goal to improve how concussion education is delivered as part of the health curriculum.

BOISE St Luke's

Resources

pedsconcussion.com




PedsConcussion
— LIVING GUIDELINE FOR —
PEDIATRIC CONCUSSION

The Living Guideline for Pediatric Concussion shares Up-to-date clinical practice guideline recommendations and tools for preventing, diagnosing, and managing pediatric concussion. The project team includes over 45 volunteer concussion experts from across the US and Canada who work together to review the latest evidence and update the clinical recommendations and tools as the evidence evolves. See the "What's New" tab for updates and scroll down for a full list of our clinical guidelines recommendations, tools, and clinical algorithms.

[Updated Concussion Handouts](#) [Living Guideline Evidence Map](#) [Download Guideline Citation](#)

Concussion Recognition, Diagnosis and Initial Medical Assessment, & Return to School





CDC HEADS UP
SAFE BRAIN. STRONGER FUTURE.

**A FACT SHEET FOR
School Nurses**

This fact sheet will help school nurses protect students from concussion or other serious brain injury; know how to recognize a concussion and what to do if one occurs; support students who return to school while recovering from a concussion; and provide steps to prevent concussion in school.

cdc.gov/heads-up



**RETURNING TO SCHOOL
AFTER A CONCUSSION**
A Fact Sheet for School Professionals

CDC HEADS UP
SAFE BRAIN. STRONGER FUTURE.

What is a Concussion?
A concussion is a type of traumatic brain injury (TBI) that results from a

What role do I play in helping a student return to school after a concussion?
Each year hundreds of thousands of K-12 students sustain a concussion as a result of a fall, motor-vehicle crash, collision on the playground or sports field.



Key Points

- Utilize updated concussion information based on current research.
- Educate your school team before concussion happens by developing a concussion management plan.
- Ongoing **team** communication – student & family, academic team and medical team.



Thank you!

Contact: 208-381-2665

parduek@slhs.org



References

- Centers for Disease Control & Prevention (2024). Head Up: Returning to school after a concussion. <https://www.cdc.gov/heads-up/guidelines/returning-to-school.html>
- Gioia, G.A. Medical-school partnership in guiding return to school following mild traumatic brain injury in youth. *Journal of Child Neurology* (2016); 31, (1):93-108.
- Kemp, A.M., O'Brien, K.H., Critical elements of return to learn for students with concussions: a scoping review. *J. Head Trauma Rehabilitation* (2021); 37,(2): E113-E128.
- Parachute (2024). Concussion. <https://parachute.ca/en/injury-topic/concussion/>
- Patricios, J. S., Schneider, K. J., Dvorak, J., Ahmed, O. H., Blauwet, C., Cantu, R. C., Davis, G. A., Echemendia, R. J., Makdissi, M., McNamee, M., Broglio, S., Emery, C. A., Feddermann-Demont, N., Fuller, G. W., Giza, C. C., Guskiewicz, K. M., Hainline, B., Iverson, G. L., Kutcher, J. S., ... Meeuwisse, W. (2023). Consensus statement on concussion in sport: The 6th international conference on concussion in sport – Amsterdam, October 2022. *British Journal of Sports Medicine*, 57, 695-711. <https://doi.org/10.1136/bjsports-2023-106898>.
- Rigney, G.H., Jo, J., Burns, C, Williams, K.L., Terry, D.P, Zuckerman, S.L. Do academic accommodations help students recover following sport-related concussion? A retrospective study of 96 athletes. *J Neurosurg Pediatrics*, (2024); (33):109-117.

